

This page to be completed by:  
Program Staff and Parent or Guardian

## Allergy Care Plan Request Form

Child's name: \_\_\_\_\_

Child's date of birth: \_\_\_\_\_

Early Learning or Child Care Program Director: Mrs. Vanshika Malhotra

Early Learning or Child Care Program: Bellevue Children's Academy - Satellite 1 & Satellite 2

Mailing Address: Bellevue Children's Academy, Satellite 1, 14719 NE 29th Pl, Bellevue, WA 98007  
Bellevue Children's Academy, Satellite 2, 14673 NE 29th Pl., Bellevue, WA 98007

Phone Number: 425-649-0791 Option 3 (Satellite 1); Option 4 (Satellite 2)

Email: preprimary@bcacademy.com

**Healthcare Provider:** The child listed above attends our program. This packet includes forms to help meet our licensing standards (WAC 110-300-0215 and 110-300-0300). **Please complete pages 2-3 and write in appropriate food substitutions in the "Additional Instructions" section on page 3.** A healthcare provider is required to provide this information and sign these forms.

**If the child has a diagnosed food intolerance, please contact the program listed above to request the Food Intolerance Care Plan Packet.**

By signing below, I give permission to my child's healthcare provider to release the health information requested in the following care plan to my child's program.

Parent or Guardian Name (Printed): \_\_\_\_\_

Parent or Guardian Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Parent or Guardian Phone Number: \_\_\_\_\_

# Allergy and Anaphylaxis Emergency Plan

American Academy of Pediatrics


DEDICATED TO THE HEALTH OF ALL CHILDREN®



Child's name: \_\_\_\_\_ Date of plan: \_\_\_\_\_

Date of birth: \_\_\_\_/\_\_\_\_/\_\_\_\_ Age \_\_\_\_ Weight: \_\_\_\_\_kg

Child has allergy to \_\_\_\_\_



- Child has asthma.  Yes  No (If yes, higher chance severe reaction)  
 Child has had anaphylaxis.  Yes  No  
 Child may carry medicine.  Yes  No  
 Child may give him/herself medicine.  Yes  No (If child refuses/is unable to self-treat, an adult must give medicine)

## IMPORTANT REMINDER

**Anaphylaxis is a potentially life-threatening, severe allergic reaction. If in doubt, give epinephrine.**

### For Severe Allergy and Anaphylaxis What to look for



If child has ANY of these severe symptoms after eating the food or having a sting, **give epinephrine.**

- Shortness of breath, wheezing, or coughing
- Skin color is pale or has a bluish color
- Weak pulse
- Fainting or dizziness
- Tight or hoarse throat
- Trouble breathing or swallowing
- Swelling of lips or tongue that bother breathing
- Vomiting or diarrhea (if severe or combined with other symptoms)
- Many hives or redness over body
- Feeling of "doom," confusion, altered consciousness, or agitation

**SPECIAL SITUATION:** If this box is checked, child has an extremely severe allergy to an insect sting or the following food(s): \_\_\_\_\_. Even if child has MILD symptoms after a sting or eating these foods, **give epinephrine.**

### Give epinephrine! What to do

1. Inject epinephrine right away! Note time when epinephrine was given.
2. Call 911.
  - Ask for ambulance with epinephrine.
  - Tell rescue squad when epinephrine was given.
3. Stay with child and:
  - Call parents and child's doctor.
  - Give a second dose of epinephrine, if symptoms get worse, continue, or do not get better in 5 minutes.
  - Keep child lying on back. If the child vomits or has trouble breathing, keep child lying on his or her side.
4. Give other medicine, if prescribed. Do not use other medicine in place of epinephrine.
  - Antihistamine
  - Inhaler/bronchodilator

### For Mild Allergic Reaction What to look for



If child has had any mild symptoms, **monitor child.**

Symptoms may include:

- Itchy nose, sneezing, itchy mouth
- A few hives
- Mild stomach nausea or discomfort

### Monitor child What to do

Stay with child and:

- Watch child closely.
- Give antihistamine (if prescribed).
- Call parents and child's doctor.
- If more than 1 symptom or symptoms of severe allergy/anaphylaxis develop, use epinephrine. (See "For Severe Allergy and Anaphylaxis.")

## Medicines/Doses

Epinephrine, intramuscular (list type): \_\_\_\_\_ Dose:  0.10 mg (7.5 kg to less than 13 kg)\*  
 0.15 mg (13 kg to less than 25 kg)  
 0.30 mg (25 kg or more)

Antihistamine, by mouth (type and dose): \_\_\_\_\_ (\*Use 0.15 mg, if 0.10 mg is not available)

Other (for example, inhaler/bronchodilator if child has asthma): \_\_\_\_\_

Parent/Guardian Authorization Signature

Date

Physician/HCP Authorization Signature

Date

# Allergy and Anaphylaxis Emergency Plan

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



Child's name: \_\_\_\_\_ Date of plan: \_\_\_\_\_

## Additional Instructions:

## Contacts

Call 911 / Rescue squad: \_\_\_\_\_

Doctor: \_\_\_\_\_ Phone: \_\_\_\_\_

Parent/Guardian: \_\_\_\_\_ Phone: \_\_\_\_\_

Parent/Guardian: \_\_\_\_\_ Phone: \_\_\_\_\_

## Other Emergency Contacts

Name/Relationship: \_\_\_\_\_ Phone: \_\_\_\_\_


Name/Relationship: \_\_\_\_\_ Phone: \_\_\_\_\_

There are many different types of Epinephrine auto-injectors. **Always follow the instructions on the medication label, as well as any child-specific instructions from parents or guardians.**

Below are common types of epinephrine auto-injectors and how to use them. **These instructions\* DO NOT replace staff training by the parent or guardian.**


## HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




## HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



## HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALCLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.



## HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



## HOW TO USE SYMJEPI™ (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPI by finger grips only and slowly insert the needle into the thigh. SYMJEPI can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



## ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

This page to be completed by:  
Program Staff and Parent or Guardian

## Additional Requirements for Care Plans

Child's name: \_\_\_\_\_

**Program Staff and Parent or Guardian:** The WAC requires that all care plans include the potential side effects and expiration date of medications. If this is not included in the care plan, write them in the table below. **You may find this information on the medication packaging or label.**

Medication Name	Expiration Date	Potential Side Effects

**Program Staff and Parent or Guardian:** The WAC requires a parent, guardian, or appointed designee to provide training to program staff about medication administration or special medical procedures listed in the child's care plan. **Use the space below to document this training.**

Employee Training Record				
Date of Training	Employee Name (Printed)	Employee Signature	Trainer Name (Printed)	Trainer Signature

**Program Staff and Parent or Guardian:** The WAC requires written consent from a child's parent or guardian before a program can administer any medications or follow a care plan that is completed by a healthcare provider. **Please have the parent or guardian sign below.**

By signing below, I give the program permission to follow this care plan as ordered by the healthcare provider.

Parent or Guardian Name (Printed): \_\_\_\_\_

Parent or Guardian Signature: \_\_\_\_\_

Date: \_\_\_\_\_

This page to be completed by:  
Parent or Guardian

## Emergency Contact Information

Child's name: \_\_\_\_\_

**Parent or Guardian:** If your child has a medical emergency, program staff need to be able to contact you or another emergency contact as quickly as possible. Please complete the following:

### Emergency Contact #1

Name: \_\_\_\_\_

Relationship to Child: \_\_\_\_\_

Phone Number: \_\_\_\_\_

### Emergency Contact #2

Name: \_\_\_\_\_

Relationship to Child: \_\_\_\_\_

Phone Number: \_\_\_\_\_

### Emergency Contact #3

Name: \_\_\_\_\_

Relationship to Child: \_\_\_\_\_

Phone Number: \_\_\_\_\_

This page to be completed by:  
Program Staff

# Medication Log

**Program Staff:** Please print a Medication Log for each medication (including any 3-Day Critical Medication).

**Child's name:** \_\_\_\_\_

**Child's date of birth:** \_\_\_\_\_

**Name of medication:** \_\_\_\_\_

Date	Time	Dose	Person Giving Medication (*Initials)	Reason Medication Was Not Given	Observed Side Effects

Initials*	Printed Name and Signature of Person Giving Medications