

This page to be completed by:
Program Staff and Parent or Guardian

Allergy Care Plan Request Form

Child's name: _____

Child's date of birth: _____

Early Learning or Child Care Program Director: Abby Maclean

Early Learning or Child Care Program: Bellevue Children's Academy - EARLY CHILDHOOD

Bellevue Children's Academy, North Campus - Satellite 1, 14719 NE 29th Pl., Bellevue, WA 98007

Mailing Address: Bellevue Children's Academy, North Campus - Satellite 2, 14673 NE 29th Pl., Bellevue, WA 98007

Phone Number: 425-649-0791 - Satellite 1 (Opt. 3); Satellite 2 (Opt. 4)

Fax Number: Satellite@bcacademy.com

Healthcare Provider: The child listed above attends our program. This packet includes forms to help meet our licensing standards for medications and individual care plans. **Please complete pages 2-3 and write in appropriate food substitutions in the "Additional Instructions" section on page 3.** A healthcare provider is required to provide this information and sign these forms.

If the child has a diagnosed food intolerance, please contact the program listed above to request the Food Intolerance Care Plan Packet.

By signing below, I give permission to my child's healthcare provider to release the health information requested in the following care plan to my child's program.

Parent or Guardian Name (Printed): _____

Parent or Guardian Signature: _____

Date: _____

Parent or Guardian Phone Number: _____

Allergy and Anaphylaxis Emergency Plan

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



Child's name: _____ Date of plan: _____

Date of birth: ____/____/____ Age ____ Weight: _____kg

Child has allergy to _____

- Child has asthma. ☐ Yes ☐ No (If yes, higher chance severe reaction)
- Child has had anaphylaxis. ☐ Yes ☐ No
- Child may carry medicine. ☐ Yes ☐ No
- Child may give him/herself medicine. ☐ Yes ☐ No (If child refuses/is unable to self-treat, an adult must give medicine)

Attach
child's
photo

IMPORTANT REMINDER

Anaphylaxis is a potentially life-threatening, severe allergic reaction. If in doubt, give epinephrine.

For Severe Allergy and Anaphylaxis What to look for



If child has ANY of these severe symptoms after eating the food or having a sting, **give epinephrine**.

- Shortness of breath, wheezing, or coughing
- Skin color is pale or has a bluish color
- Weak pulse
- Fainting or dizziness
- Tight or hoarse throat
- Trouble breathing or swallowing
- Swelling of lips or tongue that bother breathing
- Vomiting or diarrhea (if severe or combined with other symptoms)
- Many hives or redness over body
- Feeling of "doom," confusion, altered consciousness, or agitation

☐ **SPECIAL SITUATION:** If this box is checked, child has an extremely severe allergy to an insect sting or the following food(s): _____. Even if child has MILD symptoms after a sting or eating these foods, **give epinephrine**.

Give epinephrine! What to do

1. Give epinephrine right away! Note time when epinephrine was given.
2. Call 911.
 - Ask for ambulance with epinephrine.
 - Tell rescue squad when epinephrine was given.
3. Stay with child and:
 - Call parents and child's doctor.
 - Give a second dose of epinephrine, if symptoms get worse, continue, or do not get better in 5 minutes.
 - Keep child lying on back. If the child vomits or has trouble breathing, keep child lying on his or her side.
4. Give other medicine, if prescribed. Do not use other medicine in place of epinephrine.
 - Antihistamine
 - Inhaler/bronchodilator

For Mild Allergic Reaction What to look for



If child has had any mild symptoms, **monitor child**.

Symptoms may include:

- Itchy nose, sneezing, itchy mouth
- A few hives
- Mild stomach nausea or discomfort

Monitor child What to do

Stay with child and:

- Watch child closely.
- Give antihistamine (if prescribed).
- Call parents and child's doctor.
- If more than 1 symptom or symptoms of severe allergy/anaphylaxis develop, use epinephrine. (See "For Severe Allergy and Anaphylaxis.")

Medicines/Doses

Epinephrine (list type): _____ Intramuscular: ☐ 0.10 mg (7.5 kg to less than 13 kg)*
☐ 0.15 mg (13 kg to less than 25 kg)
☐ 0.30 mg (25 kg or more)
(*Use 0.15 mg, if 0.10 mg is not available)

Intranasal: ☐ 2 mg (30 kg or more)

**If more than one epinephrine is selected, then either one can be used

Antihistamine, by mouth (type and dose): _____

Other (for example, inhaler/bronchodilator if child has asthma): _____

Parent/Guardian Authorization Signature

Date

Physician/HCP Authorization Signature

Date

© 2017 American Academy of Pediatrics, Updated 10/2024. All rights reserved. Your child's doctor will tell you to do what's best for your child.
This information should not take the place of talking with your child's doctor. Page 1 of 2.

Allergy Care Plan Packet
Page 2 of 7

Allergy and Anaphylaxis Emergency Plan

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



Child's name: _____ Date of plan: _____

Additional Instructions:

Contacts

Call 911 / Rescue squad: _____

Doctor: _____ Phone: _____

Parent/Guardian: _____ Phone: _____

Parent/Guardian: _____ Phone: _____

Other Emergency Contacts

Name/Relationship: _____ Phone: _____


Name/Relationship: _____ Phone: _____

There are many different types of Epinephrine auto-injectors. **Always follow the instructions on the medication label, as well as any child-specific instructions from parents or guardians.**

Below are common types of epinephrine auto-injectors and how to use them. **These instructions* DO NOT replace staff training by the parent or guardian.**


HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENAClick®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE SYMJEPi™ (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPi by finger grips only and slowly insert the needle into the thigh. SYMJEPi can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

This page to be completed by:
Program Staff and Parent or Guardian

Additional Requirements for Care Plans

Child's name: _____

Program Staff and Parent or Guardian: The WAC requires that all care plans include the potential side effects and expiration date of medications. If this is not included in the care plan, write them in the table below. **You may find this information on the medication packaging or label.**

Medication Name	Expiration Date	Potential Side Effects

Program Staff and Parent or Guardian: The WAC requires a parent, guardian, or appointed designee to provide training to program staff about medication administration or special medical procedures listed in the child's care plan. **Use the space below to document this training.**

Employee Training Record				
Date of Training	Employee Name (Printed)	Employee Signature	Trainer Name (Printed)	Trainer Signature

Program Staff and Parent or Guardian: The WAC requires written consent from a child's parent or guardian before a program can administer any medications or follow a care plan that is completed by a healthcare provider. **Please have the parent or guardian sign below.**

By signing below, I give the program permission to follow this care plan as ordered by the healthcare provider.

Parent or Guardian Name (Printed): _____

Parent or Guardian Signature: _____

Date: _____

This page to be completed by:
Parent or Guardian

Emergency Contact Information

Child's name: _____

Parent or Guardian: If your child has a medical emergency, program staff need to be able to contact you or another emergency contact as quickly as possible. Please complete the following:

Emergency Contact #1

Name: _____

Relationship to Child: _____

Phone Number: _____

Emergency Contact #2

Name: _____

Relationship to Child: _____

Phone Number: _____

Emergency Contact #3

Name: _____

Relationship to Child: _____

Phone Number: _____

This page to be completed by:
Program Staff

Medication Log

Program Staff: Please print a Medication Log for each medication (including any 3-Day Critical Medication).

Child's name: _____

Child's date of birth: _____

Name of medication: _____

Date	Time	Dose	Person Giving Medication (*Initials)	Reason Medication Was Not Given	Observed Side Effects

Initials*	Printed Name and Signature of Person Giving Medications